

Relaxation of Static Problems

M. Ortiz

California Institute of Technology and
Rheinische Friedrich-Wilhelms Universität Bonn

with: Sergio Conti and Stefan Müller (uni-Bonn)

20th GAMM Seminar on Microstructures
Online on ZOOM, January 28-29, 2021



Outline

- *Static problems*: Stresses are the main unknowns, *equilibrium* and *minimum complementary energy* are the main operative principle (method of forces)
- Focus on the static problem of limit analysis at collapse for given limit surface of the material
- Existence, attainment, *quasiconvexity*, relaxation?
- Fused silica glass as an example of a class of materials that lack quasiconvexity and undergo relaxation through stress patterning
- Explicit constructions and numerical experiments...



Kinematic and static problems

- *Hellinger-Reissner functional:*

$$J(u, \sigma) = \int_{\Omega} \left(\sigma \cdot \varepsilon(u) - \chi(\sigma) - f \cdot u \right) dx \\ - \int_{\Gamma_D} \sigma \nu \cdot (u - g) d\mathcal{H}^{d-1} - \int_{\Gamma_N} h \cdot u d\mathcal{H}^{d-1}$$

- *Kinematic problem:* $\inf_u \sup_{\sigma} J(u, \sigma)$

- Convexity, $W = \chi^* \Rightarrow$ *Minimum potential energy*

$$\inf_u \left\{ \int_{\Omega} \left(W(\varepsilon(u)) - f \cdot u \right) dx - \int_{\Gamma_N} h \cdot u d\mathcal{H}^{d-1}, \quad u = g \text{ on } \Gamma_D \right\}$$

- Lower-semicontinuity \sim *(Morrey) quasiconvexity*



Equilibrium satisfied in limit along min sequences

Kinematic and static problems

- *Static problem:* $\sup_{\sigma} \inf_u J(u, \sigma)$
- Equivalent formulation: $u \sim$ Lagrange multipliers,

$$\sup_{\sigma} \left\{ \int_{\Gamma_D} \sigma \nu \cdot g \, d\mathcal{H}^{d-1} - \int_{\Omega} \chi(\sigma) \, dx \right\}$$


subject to: i) $\operatorname{div} \sigma + f = 0$, in Ω ;

ii) $\sigma \nu = h$, on Γ_N .

- Lower-semicontinuity \sim *sym-div quasiconvexity!*
- Primary unknown: *Stress field*
- *Equilibrium* built into solution space *ab initio*
- Complementary energy *not necessarily convex!*
- *Relaxation?* (in stress space!)



Application: Collapse of solids & structures




A steel arch bridge collapses in Yilan County, northeastern Taiwan, killing six people, on Oct. 1, 2019


Michael Ortiz
GAMM 01/21

Limit analysis at collapse

- *Stages of hardening*: Initial yield, power-law hardening and saturation



Typical stress-strain curve of elastic-plastic material showing transition from initial yield to saturation



Evolution of elastic domain from initial yield surface to *limiting surface* at saturation



Limit analysis at collapse

- *Definition of collapse:* Unconstrained incremental displacements (or velocities) at constant load

Theorem. Assume elastic-plastic solid, convex elastic domain, rate-independent behavior. Then,

$$\dot{f} = 0, \quad \dot{g} = 0, \quad \dot{h} = 0 \Rightarrow \dot{\sigma} = 0, \quad \dot{\varepsilon}^e = 0, \quad \dot{\varepsilon} = \dot{\varepsilon}^p.$$

- No incremental hardening (stress on *limit surface*)
- Behavior at collapse is *rigid-ideally plastic*
- Collapse signals the failure of *safe-load condition* and the *loss of coercivity* of rate functional
- *Static problem* for rate-independent rigid-plasticity



Limit analysis at collapse

- Recall, *general static problem*:

$$\sup_{\sigma} \left\{ \int_{\Gamma_D} \sigma \nu \cdot g \, d\mathcal{H}^{d-1} - \int_{\Omega} \chi(\sigma) \, dx \right\}$$

subject to: i) $\operatorname{div} \sigma + f = 0$, in Ω ;

ii) $\sigma \nu = h$, on Γ_N .

- Collapse*, limit surface K : $\chi(\sigma) = \begin{cases} 0, & \text{if } \sigma \in K, \\ +\infty, & \text{otherwise.} \end{cases}$
- Static problem at collapse*:

$$\sup_{\sigma} \left\{ \int_{\Gamma_D} \sigma \nu \cdot g \, d\mathcal{H}^{d-1} : \operatorname{div} \sigma = 0, \sigma(x) \in K \right\}$$

- If K convex: Equivalent *kinematic problem*



Example: Fused silica glass


Starting structure: β -cristobalite

β -cristobalite: Polymorph characterized by corner-bonded SiO_4 tetrahedra


Amorphous structure of fused silica: Obtained through the **fast quenching** of a melt

Steps taken during **quenching** process:

- Uniform temperature decrease from 5000 K to 300 K, decreasing the temperature with steps of 500 K
- Total cooling time: 470 ps



Ideal structure of β -cristobalite (adapted from <https://en.wikipedia.org/wiki/Cristobalite>)






W. Schill and S. Heyden and S. Conti and M. Ortiz, *JMPS*, **113** (2018) 105-125.

Michael Ortiz
GAMM 01/21

Example: Fused silica glass

- *Pressure-dependent plasticity*: Critical states (yielding at constant volume) define Critical State Line (CSL)
- CSL bounds the limit domain K




Limit domain K strongly non convex!

Example: Fused silica glass

Effect of a Coordination Change on the Strength of Amorphous SiO_2

CHARLES MEADE AND RAYMOND JEANLOZ

Fig. 1. Maximum shear stress in silica glass at room temperature and average pressures (\bar{P}) between 8.6 and 81 GPa. Each point corresponds to a separate sample, and the heavy line shows the general trend of the data. The shear stress is determined from Eq. 1, and it is a measure of the yield strength of the sample at high pressures. The error bars represent the combined uncertainties from the measurements of h and $\partial P/\partial r$. The open circles show the strength of samples that were initially compressed to 50 GPa, unloaded, and then recompressed. The arrow marks the zero pressure strength of silica glass (19).



1072

SCIENCE, VOL. 241

Anomalous shear strength of fused silica glass is well-documented in the geophysics literature

Michael Ortiz
GAMM 01/21



Symmetric div-quasiconvexity

- Recall, static problem at collapse:

$$\sup_{\sigma} \left\{ \int_{\Gamma_D} \sigma \nu \cdot g \, d\mathcal{H}^{d-1} : \operatorname{div} \sigma = 0, \sigma(x) \in K \right\}$$



- If K is *strongly non-convex*, non-attainment.
- What is an appropriate notion of *quasiconvexity*, *relaxation*, *envelops*, *constructions*?
- Symmetric div-quasiconvexity* (sdqc), a special case of *\mathcal{A} -quasiconvexity* (Fonseca-Müller, 1999), implies *existence of solutions* of the static problem
- Relaxed problem defined by *sdqc envelop* K^{sdqc}
- Exact constructions* for isotropic domains...



Symmetric div-quasiconvexity

Div-quasiconvex envelop of glass elastic domain:


- **Theorem** (Tartar'85). *The function $f(\sigma) = 2|\sigma|^2 - \text{tr}(\sigma)^2$ is div-quasiconvex.*
- **Theorem.** *The set $\{\sigma : q^2 \leq c^2 + \frac{3}{4}(p - p_0)^2\}$ is sym div-quasiconvex.*
- **Theorem.** *The sym div-quasiconvex envelop K^{sdqc} of $K(\text{Si}_2\text{O})$ is:*



S. Conti, S. Muller and M. Ortiz, *ARMA*, **235** (2020) 841-880.

Michael Ortiz
GAMM 01/21

Symmetric div-quasiconvexity



W. Schill, S. Heyden, S. Conti and M. Ortiz, *JMPS*, **113** (2018) 105-125.

Michael Ortiz
GAMM 01/21

Concluding Remarks

- Static problems lacking attainment have been studied relatively less than kinematic problems
- Symmetric div-quasiconvexity provides a practical and compelling illustration of the abstract concept of \mathcal{A} -quasiconvexity of Fonseca and Müller
- Static problems set forth a non-standard type of relaxation achieved by means of equilibrated fine-oscillations in stress space
- The isotropic case is amenable to explicit envelop constructions that characterize rigorously and efficiently the macroscopic effective behavior of large classes of materials...



Can we prevent collapse?



